

# Social Psychiatry and Psychiatric Epidemiology

## High prevalence of psychiatric disorders in Iran: results from the Iranian Mental Health Survey, 2011 --Manuscript Draft--

<b>Manuscript Number:</b>	SPPE-D-14-00462
<b>Full Title:</b>	High prevalence of psychiatric disorders in Iran: results from the Iranian Mental Health Survey, 2011
<b>Article Type:</b>	Original Paper
<b>Keywords:</b>	psychiatric disorders; Epidemiology; Prevalence; Iran; Mental Health Survey
<b>Abstract:</b>	<p><b>Purpose:</b> The study aimed to estimate the 12-month prevalence, severity and correlates of psychiatric disorders in Iranian adult population.</p> <p><b>Methods:</b> The Iran Mental Health Survey (IranMHS) was conducted through a multistage cluster sampling design. A total of 7886 community dwelling residents aged 15-64 were recruited. 12-month diagnoses of DSM-IV psychiatric disorders using a validated Persian translation of the Composite International Diagnostic Interview (CIDI; version 2.1). The Structured Clinical Interview for DSM-IV Axis I disorders (SCID-I) was administered by a psychiatrist on subjects screened positive for the presence of a psychotic disorder. Severity of psychiatric disorders was determined using criteria that included markers for disability, particular diagnoses and suicide attempts.</p> <p><b>Results:</b> The response rate was 86.2%. The 12-month weighted prevalence of "any psychiatric disorders" was 23.6% [95% confidence interval (CI): 22.4-24.8] with 26.5% of women and 20.8% of men having one or more psychiatric disorders. The most common category of psychiatric disorders was any anxiety disorder (15.6%) and the most prevalent particular disorder was major depressive disorder (12.7%). A 12-month psychotic disorder was observed in 0.5% of the population (95% CI: 0.33-0.66). Almost two-thirds (63.8%) of individuals with a mental disorder had moderate or serious illness. Unemployment, being widowed/divorced and urban living were associated with a greater likelihood of 12-month disorders; while, higher socioeconomic status and having a university degree were associated with a lower likelihood.</p> <p><b>Conclusion:</b> The high prevalence of psychiatric disorders and particularly major depression merits further attention in the country's mental health policy and program planning.</p>

## RESEARCH REPORT

### **High prevalence of psychiatric disorders in Iran: results from the Iranian Mental Health Survey, 2011**

#### **Vandad Sharifi, MD, MPH**

Department of Psychiatry and Psychiatry and Psychology Research Center, Tehran University of Medical Sciences, Tehran, Iran, Email address: [vsharifi@tums.ac.ir](mailto:vsharifi@tums.ac.ir)

#### **Masoumeh Amin-Esmaeili\*, MD, MPH (Corresponding author)**

Department for Mental Health and Substance Use, Iranian Research Center for HIV/AIDS (IRCHA) and Iranian National Center for Addiction Studies (INCAS), Iranian Institute for Reduction of High-Risk Behaviors, Tehran University of Medical Sciences, Tehran, Iran, Email address: [m.aminesmaeeli@farabi.tums.ac.ir](mailto:m.aminesmaeeli@farabi.tums.ac.ir)

#### **Ahmad Hajebi\*, MD (Corresponding author)**

Mental Health Research Center, Tehran Institute of Psychiatry- Faculty of Behavioral Sciences and Mental Health, Iran University of Medical Sciences, Tehran, Iran, Email Address: [hajebi.a@iums.ac.ir](mailto:hajebi.a@iums.ac.ir)

#### **Abbas Motevalian, MD, PHD**

Department of Epidemiology, School of Public Health, Iran University of Medical Sciences, Tehran, Iran, Email address: [amotevalian@yahoo.com](mailto:amotevalian@yahoo.com)

#### **Reza Radgoodarzi, MD**

Iranian National Center for Addiction Studies (INCAS), Iranian Institute for Reduction of High-Risk Behaviors, Tehran University of Medical Sciences and Tous Clinic, Tehran, Iran, Email address: [rezaradus@gmail.com](mailto:rezaradus@gmail.com)

#### **Mitra Hefazi, MD**

Department for Mental Health and Substance Use, Iranian Research Center for HIV/AIDS (IRCHA) and Iranian National Center for Addiction Studies (INCAS), Iranian Institute for Reduction of High-Risk Behaviors, Tehran University of Medical Sciences, Tehran, Iran, Email address: [mitra\\_hefazi@yahoo.com](mailto:mitra_hefazi@yahoo.com)

#### **Afarin Rahimi-Movaghar, MD, MPH**

Iranian National Center for Addiction Studies (INCAS), Iranian Institute for Reduction of High-Risk Behaviors, Tehran University of Medical Sciences, Tehran, Iran, Email address: [rahimia@tums.ac.ir](mailto:rahimia@tums.ac.ir)

#### **\*Joint corresponding authors:**

#### **Dr. M. Amin-Esmaeili, M.D., M.P.H.**

Address: Imam Khomeini Hospital, Keshavarz Blvd., Tehran, Iran  
Postal code: 14197-33141; Telefax: +98-21-66918899; +98-21-66947984; email: [m.aminesmaeeli@farabi.tums.ac.ir](mailto:m.aminesmaeeli@farabi.tums.ac.ir)

#### **Dr. A. Hajebi, M.D.**

1  
2  
3  
4 Address: Tehran Psychiatric Institute, Mansouri St., Nyayesh Ave., Sattarkhan Ave., Tehran, Iran; Tel:  
5 +98-21-66506862; email: [hajebi.a@iums.ac.ir](mailto:hajebi.a@iums.ac.ir)  
6

7  
8 **Authors' contributions:**

9  
10 All authors were involved in study conceptualization, design, training of the interviewers, quality control  
11 and interpretation of the results. A. Rahimi Movaghar was responsible for the overall management of the  
12 IranMHS. Statistical analyses were performed by A. Motevalian. V. Sharifi, M. Amin-Esmaili and A.  
13 Hajebi drafted the manuscript. All authors contributed to the final version of the article.  
14

15  
16 **Conflict of interest:**

17 None  
18

19  
20 **word count** : 4105  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4  
5 **ABSTRACT**  
6

7  
8 **Purpose:** The study aimed to estimate the 12-month prevalence, severity and correlates of psychiatric  
9 disorders in Iranian adult population.  
10

11  
12  
13 **Methods:**  
14

15  
16 The Iranian Mental Health Survey (IranMHS) was conducted through a multistage cluster sampling  
17 design. A total of 7886 community dwelling residents aged 15–64 were recruited. 12-month diagnoses of  
18 DSM-IV psychiatric disorders using a validated Persian translation of the Composite International  
19 Diagnostic Interview (CIDI; version 2.1). The Structured Clinical Interview for DSM-IV Axis I disorders  
20 (SCID-I) was administered by a psychiatrist on subjects screened positive for the presence of a psychotic  
21 disorder. Severity of psychiatric disorders was determined using criteria that included markers for  
22 disability, particular diagnoses and suicide attempts.  
23  
24  
25  
26  
27  
28  
29  
30  
31

32  
33 **Results:** The response rate was 86.2%. The 12-month weighted prevalence of “any psychiatric disorders”  
34 was 23.6% [95% confidence interval (CI): 22.4-24.8] with 26.5% of women and 20.8% of men having  
35 one or more psychiatric disorders. The most common category of psychiatric disorders was any anxiety  
36 disorder (15.6%) and the most prevalent particular disorder was major depressive disorder (12.7%). A 12-  
37 month psychotic disorder was observed in 0.5% of the population (95% CI: 0.33-0.66). Almost two-thirds  
38 (63.8%) of individuals with a mental disorder had moderate or serious illness. Unemployment, being  
39 widowed/divorced and urban living were associated with a greater likelihood of 12-month disorders;  
40 while, higher socioeconomic status and having a university degree were associated with a lower  
41 likelihood.  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51

52  
53  
54 **Conclusion:** The high prevalence of psychiatric disorders and particularly major depression merits further  
55 attention in the country’s mental health policy and program planning.  
56  
57  
58

59 **Key words:** psychiatric disorders; Epidemiology; Prevalence; Iran; Mental Health Survey.  
60  
61

## Introduction

Iran is a large country located in the Middle East with a population of more than 75 million, making it the 17<sup>th</sup> most populated country in the world [1]. The country has been undergoing rapid demographic, societal and economic changes in the last few decades. The urban dwelling proportion of the population has increased from 27% in 1950 to 71.4% in 2011 [1]. The population growth rate has also decreased, leading to a change in age structure, with young people aged 20-29 years becoming the largest age group in the country [1]. Iran has faced several adversities in the last few decades; including an 8-year war in the 1980s, severe economic sanctions and numerous natural disasters in more recent years. Nonetheless, during the same period, the national health status improved as evidenced by increased life expectancy, and reductions in infant and maternal mortality rates [2]. There were also advances in the health care system of the country including increased per capita expenditure on health and improved access to services [2]. Little is known about the possible impact of the above-mentioned socio-demographic, economic, and health care system developments on the population's mental health status. Addressing the mental health care needs of the population requires up-to-date national level information. The only available nationwide study of the mental health status of the Iranian population dates back to more than 10 years ago [3,4].

The Iranian Mental Health Survey (IranMHS) was designed and conducted to provide first-hand information to aid policy-making at the national level. It is a comprehensive and nationally representative survey of the prevalence and severity of psychiatric conditions, as well as the current status of service utilization and costs of mental health care. This article reports the results pertaining to the former objective, and aims to present the 12-month prevalence and severity of psychiatric disorders and correlates of the presence of psychiatric disorder in Iranian community-dwelling population.

## Methods

### *Sample*

The methodology of the IranMHS is described in detail elsewhere [5]. Briefly, this study was a household survey conducted on individuals 15-64 years-old residing in Iran. A three stage cluster sampling method was employed: First, 1525 blocks were selected proportional to the size of each province based on the latest national census database [6]. In the second stage, households in each block were counted and 6 households in each block were selected through systematic random sampling. In the third stage, the individual to be interviewed in each selected household was determined by using a method described by Kish [7]. Individuals who could not comprehend Persian (the official and widely used language in Iran) or had severe physical or mental disability which made them unable to participate in the study, and those with nationalities other than Iranian were excluded. The selected individuals who were unable to participate or refused participation were not replaced. However, measures were taken to achieve the highest response, such as repeated visits for no shows. The survey was conducted from January to June 2011 and a total of 7886 individuals participated in the survey.

### *Measures*

The 12-month version of the Composite International Diagnostic Interview (CIDI, version 2.1), was employed to determine the presence of the following psychiatric disorders according to the Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition (DSM-IV): major depressive disorder, dysthymia, bipolar I disorder, panic disorder (with and without agoraphobia), agoraphobia without panic, obsessive-compulsive disorder, social phobia, post-traumatic stress disorder, and alcohol and other substance use disorders. The substances included cannabis, opioids, amphetamine-type stimulants, hallucinogens and inhalants. Because the abuse of other substances (e.g, cocaine) is rare in Iran, we did not assess each of these "other drugs" separately. But subjects were asked if they had used any other substances and If positive relevant questions were asked for the specified drug.

1  
2  
3  
4 The CIDI is a structured interview developed by the World Health Organization (WHO) designed for use  
5  
6 by trained interviewers who maybe lay persons. This tool has shown to have high inter-rater and test-  
7  
8 retest reliability and high validity for most diagnoses in different settings and languages [8,9]. The Persian  
9  
10 translation of CIDI 2.1 [10] has been validated and has shown adequate psychometric properties; with the  
11  
12 exception of psychotic disorders, which are difficult to assess in a structured interview. Also, in a pilot  
13  
14 study in a sample from general population, we observed good inter-rater reliability of diagnoses made by  
15  
16 the Persian translation of the CIDI 2.1 (kappas of 0.5 and more for different psychiatric disorders) [5].  
17  
18  
19

20  
21 To estimate the prevalence of psychotic disorders, we adopted a two stage procedure as proposed by other  
22  
23 investigators [11]. In the first stage (screening), the participants were inquired about the presence of self-  
24  
25 reported delusional beliefs and hallucinations using the items from the G section of the lifetime version of  
26  
27 the CIDI. Subsequently, all subjects who had endorsed one or more items, as well as those with a history  
28  
29 of hospitalization for a psychiatric illness were contacted to participate in the second stage. The latter  
30  
31 incorporated telephone administration of the Persian translation of the Structured Clinical Interview for  
32  
33 DSM-IV Axis I disorders (SCID-I) [12] to determine the presence of *any primary psychotic disorder* (not  
34  
35 attributable to a substance of abuse or a general medical condition) in the past 12 months. The source of  
36  
37 information was both the subject and a significant other that usually included either a spouse or a first  
38  
39 degree relative. We did not differentiate the specific types of psychotic illnesses (such as schizophrenia or  
40  
41 affective psychoses) as we were only interested in estimating the prevalence of any illness with psychotic  
42  
43 features as an indicator of a very severe mental illness. SCID-I was administered by a psychiatrist. The  
44  
45 Persian translation of the SCID-I yields DSM-IV diagnoses with good reliability and validity [12]. The  
46  
47 sensitivity of the telephone-administered Persian version of SCID-I for psychotic disorders is comparable  
48  
49 to the sensitivity of the face-to-face administered version of the instrument [13].  
50  
51  
52  
53  
54

55 Sheehan Disability Scale (SDS) [14] was used to assess the degree of disability associated with the  
56  
57 presence of any mental illness. SDS is widely used in general population surveys [15-17] to assess  
58  
59 disability in work role performance, household maintenance, social life, and intimate relationships.  
60  
61

1  
2  
3  
4 Disability on each domain is rated on a 0-10 visual analogue scales with possible ratings of none (0), mild  
5  
6 (1-3), moderate (4-6), and severe disability (7-10) [14]. Similar to the English version, the Persian  
7  
8 translation of SDS has been shown to be reliable and valid [18]. Days out of role because of a psychiatric  
9  
10 disorder was also assessed as a measure of disability and severity (see below). Participants were asked  
11  
12 about the number of days in the past 12 months in which they were totally unable to carry out their  
13  
14 normal daily activities due to any psychiatric condition.  
15  
16

17  
18 Based on psychiatric diagnoses, symptoms and disability ratings, participants were classified into those  
19  
20 with serious, moderate or mild psychiatric disorders according to the criteria used in the World Mental  
21  
22 Health survey [15] with some minor modifications. Those with a serious disorder had either a psychotic  
23  
24 disorder, bipolar I disorder or substance dependence with a physiological dependence syndrome; had  
25  
26 made a suicide attempt in the presence of any other DSM-IV disorder; or had reported severe role  
27  
28 impairment due to a mental disorder in at least two areas on the SDS. Subjects not classified as having a  
29  
30 serious disorder were classified as moderate if interference was rated as at least moderate in at least two  
31  
32 SDS domains, or if the subject had substance dependence without a physiological dependence syndrome.  
33  
34 All other participants were classified as mild.  
35  
36  
37

38  
39 Socioeconomic status of participants was assessed using a questionnaire which has been used in a  
40  
41 previous nationwide survey [19]. The questionnaire ascertains the status based on household assets, such  
42  
43 as accommodation, accessories, and possession of cars [20].  
44  
45

#### 46 *Interviewer training and procedures*

47

48  
49 Interviewers were comprised of 232 professionals with master's degrees in psychology or a BA degree  
50  
51 with a psychology major. Training consisted of a 7-day, 56-hours workshop of instruction and practice  
52  
53 sessions that included in-depth training on administration of CIDI using the standard protocols and  
54  
55 training materials developed by the World Health Organization [21,22]. Each Interviewer also took part in  
56  
57 at least 2 pilot interviews with real subjects before the main phase of the study; and received appropriate  
58  
59  
60  
61  
62  
63  
64  
65



1  
2  
3  
4 feedback from experienced interviewers. In each province, a local executive manager and a supervisor (a  
5  
6 member of research team) were assigned to ensure the quality of the procedures.  
7  
8

9  
10 The feasibility and acceptability of the interviews and the limitations were assessed in a pilot study in a  
11 selected area of the capital, Tehran, as well as an urban and a rural area of the Khoozestan province in  
12 South-west of Iran. We modified the data collection and quality control procedures for the main study  
13  
14 after the feedback from the implementation of the pilot phase.  
15  
16  
17  
18

19 All interviews were conducted face-to-face at subjects' place of residence. All completed questionnaires  
20 were reviewed and edited by the local executive manager. A third of all questionnaires were double-  
21 checked by a supervisor. Errors or inconsistencies were returned to the interviewers for resolution. This  
22  
23 involved re-interview with the participants in some cases. The research protocol was approved by the  
24  
25 Ethics Committee of Tehran University of Medical Sciences in Iran. Informed consents were obtained  
26  
27 from all participants and all data were processed anonymously.  
28  
29  
30  
31  
32

### 33 *Statistical Analysis*

34  
35  
36 Analyses were conducted in two stages. First, we computed prevalence estimates and 95% confidence  
37 intervals for individual DSM-IV disorders as well as for any 12-month psychiatric disorder. Next, we  
38 used logistic regression models to examine the correlates of psychiatric disorders in this setting. Survey  
39 weights were used in all analyses. The consolidated weights were the joint product of inverse probability  
40  
41 of unit selection into the sample, non-response weights and post-stratification weights. To calculate the  
42  
43 post-stratification weights, the proportion of subjects in each stratum in National Census 2006 was  
44  
45 divided by the proportion of the same group in the sample. Based on 5-year age groups, sex and  
46  
47 urbanicity status in each of 31 country provinces; 1240 post-stratification weights were generated.  
48  
49

50 Among 475 subjects who were screened positive for psychotic disorders, 118 did not respond to the  
51 telephone SCID interview. We regarded them as item non-responses and used the method of multiple  
52 imputations to impute missing responses [23]. The model for imputation consisted of a logistic regression  
53  
54  
55  
56  
57  
58  
59  
60  
61

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

to predict SCID 12-month diagnoses of psychotic disorders from the screening questions in the baseline survey along with demographic, service use and disability data (AUC=0.96). The regression was weighted to account for sampling probabilities of selection.

Analyses were conducted using STATA statistical package, version 12 (STATA Corporation, College Station, TX, USA, 2009). All percentages reported are weighted unless stated otherwise.

1  
2  
3  
4  
5  
6  
7 **Results**  
8

9 Of 9148 individuals approached by interviewers, 7886 (86.2%) agreed to participate in the study. The  
10 non-responder group included a higher percentage of men (56.3% vs 43.0%,  $p < 0.001$ ) and were slightly  
11 older (34.7 vs 33.0,  $p = 0.004$ ). The characteristics of the samples are presented in Table 1.  
12  
13  
14

15 -----Insert Table 1 about here-----  
16  
17

18  
19 *Twelve-month prevalence and severity of psychiatric disorders*  
20

21  
22 The 12-month prevalence of psychiatric disorders is presented in Table 2. As shown, 23.6% of the Iranian  
23 population aged 15 to 64 years met the criteria for at least one DSM-IV psychiatric disorder in the past 12  
24 months. In total, 1067 (13.4%) met the criteria for only one disorder, 493 (6.3%) two and 328 (3.9%)  
25 three or more disorders.  
26  
27  
28  
29

30  
31  
32 The most prevalent group of disorders was the group of anxiety disorders (15.6%), followed by mood  
33 disorders (14.6%). The most prevalent particular DSM-IV disorder was major depressive disorder  
34 (12.7%), followed by generalized anxiety disorder (5.2%) and obsessive-compulsive disorder (5.1%). The  
35 prevalence estimates were significantly higher among women than men for certain psychiatric disorders  
36 including major depressive disorder, agoraphobia without panic, social phobia and obsessive-compulsive  
37 disorder (Table 2). The characteristics of participants according to groups of psychiatric disorders are  
38 presented in online resource 1.  
39  
40  
41  
42  
43  
44  
45  
46  
47

48 -----Insert Table 2 about here-----  
49  
50

51 Most psychiatric disorders split almost evenly in terms of level of severity (see Table 3). We compared  
52 mean days out of role for external validation of our severity criteria. For the no disorder category, mean  
53 days out of role was 2.6 (95% CI: 2.3-3.0). The values for the mild, moderate and serious groups were 8.3  
54 (6.1-10.5), 17.1 (13.4-20.7), and 35.6 (30.1-41.2) days, respectively. This finding supports the validity of  
55 the SDS ratings.  
56  
57  
58  
59  
60  
61

-----Insert Tables3 about here-----

*Socio-demographic correlates of any psychiatric disorder*

Results of unadjusted and adjusted logistic regression analyses of the socio-demographic correlates of *any psychiatric disorder* are presented in Table 4. Women had a greater odds of meeting the criteria for any psychiatric disorder [Adjusted Odds Ratio (aOR)=1.47], as did previously married participants compared to those who were never married (aOR=1.77), and the unemployed (aOR=1.81). In contrast, individuals with a university education had a lower odds of meeting the criteria for a psychiatric disorder compared to illiterate individuals (aOR=0.65), as did those residing in rural areas compared to city-dwellers (aOR=0.71), and individuals in middle or upper socio-economic classes compared to the lower class (aOR=0.78 and 0.65, respectively).

-----Insert Table 4 about here-----

**Discussion**

The IranMHS aimed to build upon and improve the methodology of previous national surveys in the country [3,4]. We used a reliable and valid translation of an internationally well-known and widely-used instrument (i.e., CIDI) for ascertainment of psychiatric disorders that has been widely used in numerous international studies. The reliability and validity of the locally-adapted instrument was rigorously assessed [10]. Further, we employed local mental health professionals with clinical experience instead of lay interviewers. We designed and implemented a strict and comprehensive quality assurance protocol. Unlike some previous surveys in this setting, in which participants were selected for full interview based on the results of a screening, all participants received the full interview. The response rate was high enough to minimize bias.

The results of the survey suggest that almost one out of every four Iranians in the 15-64 years age range experiences one or more psychiatric disorders in a 12-month period and almost one out of every seven, experience a psychiatric disorder associated with severe to moderate disability. The estimate is one of the

1  
2  
3  
4 highest compared with some other nationwide studies using this or other versions of the CIDI. For  
5  
6 example, the World Mental Health (WMH) surveys conducted in several European, African and Asian  
7  
8 countries in 2001-2003 reported estimates varying between 4.3% in Shanghai, China to 26.4% in the US,  
9  
10 with the interquartile range (IQR) of 9.1% to 16.9% [15]. Among studies using CIDI versions 2.1 and  
11  
12 older, Bijl and Ravelli (2000) reported a 12-month prevalence of 23.5% in Netherlands and Henderson et  
13  
14 al (2000) reported a 17.7% estimate in Australia [24,25]. Our estimate is also higher than two previous  
15  
16 national surveys and several local population-based studies in Iran [3,4,26]. However, comparison of  
17  
18 prevalence estimates from different surveys is difficult as such estimates are highly dependent on  
19  
20 sampling method, instruments used to ascertain disorders, and definition of psychiatric disorders. For  
21  
22 example, in another survey of the Iranian population, Noorbala and colleagues [3,4] reported a 21%  
23  
24 prevalence of psychiatric disorders using a screening instrument (General Health Questionnaire; GHQ-  
25  
26 28). Whereas, Mohammadi and colleagues [3,4] reported a 12-month prevalence of 10.8% using the  
27  
28 Schedule for Affective Disorders and Schizophrenia (SADS). Comparisons can only be reliably made  
29  
30 across surveys using the same sampling methods and instruments. In the absence of such surveys  
31  
32 conducted over time in this setting, we cannot examine trends in prevalence of disorders as well.  
33  
34 Nevertheless, it is noteworthy that our estimate is higher than the median prevalence of all surveys that  
35  
36 have used diagnostic instruments (18.6%) as reported in a systematic review of prevalence studies in Iran  
37  
38 [26].  
39  
40  
41  
42  
43  
44

45 The relatively high prevalence estimate of mental disorders in the country calls for immediate response  
46  
47 with regard to mental health policy and program planning in the country. Despite some progress in recent  
48  
49 decades, mental health services have not yet been optimally developed in the Iran, and there is still a lack  
50  
51 of qualified mental health professionals in the country. Based on the latest available data [27] only 1.49  
52  
53 psychiatrists and 2.19 psychologists per 100,000 population work in the mental health sector and most of  
54  
55 them work in big cities.  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4 Higher prevalence among females, the unemployed, urban residents, and those with lower socio-  
5 economic status is consistent with the result of other studies across the globe [26,28,29]. However, unlike  
6 studies in other settings [28], we did not observe a significant difference between those currently married  
7 and the never married. Our finding replicates many previous studies in Iran (e.g.,[3]).One possible  
8 explanation, is that a great majority of single individuals in the country live with their parents until getting  
9 married, and therefore may not experience the stressors of independent living. Whereas, married  
10 individuals and especially married women face a number of social and economic stressors commonly  
11 associated with psychiatric disorders.  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22

23 The current study shows that anxiety disorders comprise the most common category of mental disorders.  
24 This finding is consistent with findings from a number of other countries surveyed as part of the World  
25 Mental Health surveys (WMH) [15]. However, our estimate for this category of mental illness (15.6%)  
26 surpasses all estimates in these countries with the exception of the United States (18.2%). Other  
27 nationally representative studies in Canada and the US (NCS-R) that have used CIDI.2 version reported  
28 estimates equal to 12% and 17.2%, respectively [30,31]. If we were to add simple phobias to our  
29 assessment, then we could expect having even higher rates.  
30  
31  
32  
33  
34  
35  
36  
37  
38

39 In contrast, the prevalence of depressive disorders was markedly higher in Iran than in many other  
40 countries [32]. In the WMH surveys, the average 12-month prevalence of major depressive disorder was  
41 5.5% in 10 high income countries and 5.9% in eight low to middle income countries with the highest  
42 reported prevalence being 10.6% in Brazil [32]. A systematic review of prevalence studies carried out  
43 before 2004 in different parts of Iran estimated the pooled mean prevalence of 4.1% for studies that have  
44 used diagnostic interviews [33]. The prevalence estimate in our study was three times higher than this  
45 number and suggests that there might have been increase in prevalence of depression in the country  
46 between early 2000s and 2011 when our study was conducted. Although currently debated, [34] there is  
47 some evidence of incremental trend in depressive disorders in recent years at least in some countries (e.g.,  
48 Denmark, Australia and Canada) [35-37]. The possibility of an increasing trend in Iran is also supported  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4 by other evidence from surveys conducted in Tehran showing increasing levels of psychological distress,  
5 including depression, in recent years [38]. The age structure of the Iranian population with the  
6 preponderance of the youth can not explain this finding, as we did not observe different prevalence  
7 estimates between the young and the middle age groups. Nevertheless, this unexpectedly high prevalence  
8 estimate highlights the burden of depression in the country and demands urgent action to tackle the  
9 problem.

10  
11 Our survey was the first national household survey on substance use disorders in Iran. The estimates,  
12 however, should be viewed with caution as assessments based on self-report might grossly underestimate  
13 the extent of substance use disorder due to social stigma and legal ramifications. Nevertheless, it is  
14 noteworthy that 2.8% of the population reported substance abuse or dependence, which is higher than  
15 inter quartile range of 0.8% to 2.6% from other countries in the WMH initiatives [15].

16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31 Estimating the prevalence of psychotic disorders has always been a challenging task. These illnesses are  
32 uncommon in the general population and the assessments require large sample sizes. Furthermore, relying  
33 solely on structured interviews has shortcomings, mainly because of the poor psychometric properties of  
34 these instruments. Assessments based exclusively on self-reports of the participants are prone to  
35 misinterpretation of the questions and underreporting of psychotic symptoms due to lack of insight. To  
36 overcome these barriers, we adopted a two stage procedure as proposed by other investigators [11]. The  
37 12-month prevalence of psychotic illnesses in this population was 0.50%, which is comparable to the  
38 prevalence estimate reported by Saha et al from their systematic review of previous prevalence studies  
39 [39] showing that that the median period prevalence of schizophrenia is equal to 0.33. If we had access to  
40 national health registries or databases of the institutional population, that is currently non-existent in the  
41 country, or had we interviewed a sample of screen negative subjects, then the prevalence estimate would  
42 be higher than what we obtained.

1  
2  
3  
4 Determining the severity of psychiatric disorder has important implications with regard to estimating the  
5 societal burden of these disorders and allocation of scarce health resources. Almost a third of subjects  
6 with any form of psychiatric illnesses could be allocated to each of our three categories of severity. Our  
7 findings are comparable to findings from other countries as included in the WMH surveys, considering  
8 that our severity criterion was similar to the one used in those surveys. Serious illnesses accounted for  
9 almost a third of all cases of psychiatric disorders in many countries; however, the estimates ranged from  
10 8.9% in Nigeria to 30.3% in Mexico [15].  
11  
12  
13  
14  
15  
16  
17  
18  
19

20 Any interpretation of the results of this study should be done in light of its limitations. First, our study had  
21 two major problems inherent in many cross-sectional household surveys, namely, the retrospective nature  
22 of queries and the exclusion of institutionalized or homeless people. The latter may not highly impact  
23 prevalent disorders, but it might affect less common illnesses such as psychotic and substance use  
24 disorders which are more likely to be associated with institutionalization. Second, there may have been  
25 underestimations in the prevalence of highly sensitive issues such as substance use disorders. Use and  
26 possession of alcohol and other substances are strictly punishable by law in the country and it is quite  
27 likely that a group of participants might not have reported their use. Third, CIDI is not a clinician-  
28 administered diagnostic interview and is not suitable for a comprehensive assessment required for  
29 diagnostic decision making. Nonetheless, we employed interviewer with past training in psychology and  
30 some degree of clinical experience instead of “lay interviewers”. In addition, a validation study has shown  
31 that the Persian translation administered by the same group yields reliable and valid diagnoses. Fourth,  
32 the diagnostic assessments relied on the 2<sup>nd</sup> version of the CIDI. We were unable to use the 3<sup>rd</sup> version of  
33 the CIDI (WMH-CIDI) at the time of the survey planning. However, the 2<sup>nd</sup> version is a widely used  
34 instrument in previous population surveys (e.g.,[40,41]) and we have used a reliable and valid translation  
35 of the instrument. And finally, we did not include disorders such as simple phobias, eating disorders or  
36 sleep disorders. This decision was made merely in order to shorten the interview time; bearing in mind  
37 that these disorders were assumed to impose a lower burden of disease.  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65



1  
2  
3  
4 **Conclusion**  
5

6  
7 In conclusion, the IranMHS found a relatively high prevalence of mental disorders in the country with  
8  
9 more than two thirds of people having a psychiatric disorder belonging to moderate or severe categories.

10  
11 Like many other lower and middle income countries, the share of Iranian health care budget and resources  
12  
13 that is allocated to management of these conditions is far less than would be appropriate given their social  
14  
15 burden [42]. The findings call for allocation of greater financial and human resources to the care of these  
16  
17 disorders in the country. These efforts should be coupled with increased funding for research on the  
18  
19 etiology and treatment of these conditions and efforts to improve access to care and reduce the social  
20  
21 stigma of psychiatric illness. Continued monitoring of the mental health status of the country in future  
22  
23 years could provide a gauge for the success of these efforts.  
24  
25  
26

27 **Acknowledgements**  
28

29  
30 The Iranian Mental Health Survey (IranMHS) was financially supported by the Ministry of Health  
31  
32 through the contract number 132-1491 of Tehran University of Medical Sciences and the contract number  
33  
34 89-D-432-143 of Mental Health Research Network. The funding source had no role in the study design;  
35  
36 the collection, analysis and interpretation of data; the writing of the report; and in the decision to submit  
37  
38 the article for publication.  
39  
40

41  
42 We wish to thank Dr. Ronald C. Kessler and Dr. Ramin Mojtabai for their invaluable contributions to our  
43  
44 study. We also acknowledge supports received from Dr. Hassan Emami Razavi, Dr. Ali Reza  
45  
46 Mesdaghinia, Dr. Mohammad Bagher Saberi Zafarghandi, Dr. Abbas Ali Nasehi, Dr. Mohammad Reza  
47  
48 Mohammadi and Dr. Minoos Mohraz and contributions of Dr. Samaneh Kariman, Maryam Gholamrezaei,  
49  
50 Leila Moazami Goodarzi, Farideh Kolahi Heshmat, Dr. Reza Mahdavi, Dr. Siamak Molavi and Fahima  
51  
52 Farrahi in conducting the study. We also wish to thank provincial executive managers and interviewers,  
53  
54 as well as all the participants in the study.  
55  
56

57 **Conflict of interest**  
58

59  
60 The authors have no conflicts of interests.  
61  
62

## References

1. Statistical Center of Iran (2012) Selected findings of the 2011 national population and housing census. <http://www.amar.org.ir/Portals/1/Iran/census-2.pdf>. Accessed 19 Dec 2012
2. World Health Organization (2011) Global Health Observatory (GHO), Islamic Republic of Iran: General health statistical profile. WHO. <http://www.who.int/gho/countries/irn.pdf>. Accessed 2013-06-25
3. Noorbala AA, Yazdi SAB, Yasamy MT, Mohammad K (2004) Mental health survey of the adult population in Iran. *Br J Psychiatry* 184: 70-73
4. Mohammadi MR, Davidian H, Noorbala AA et al. (2005) An epidemiological survey of psychiatric disorders in Iran. *Clin Pract Epidemiol Ment Health* 1: 16. doi:10.1186/1745-0179-1-1
5. Rahimi-Movaghar A, Amin-Esmaeili M, Sharifi V, Hajebi A, Hefazi M, Rad Goodarzi R, Motevalian SA (2014) The Iranian Mental Health Survey: design and field procedures. *Iran J Psychiatry* 9 96-109
6. Statistical Center of Iran (2006) Selected findings of the 2006 national population and housing census. <http://www.amar.org.ir/Portals/2/Files1385/kollisarshomari85.pdf>. Accessed 7 Mar 2013
7. Kish L (1949) A Procedure for Objective Respondent Selection within the Household. *Journal of the American Statistical Association* 44: 380-387
8. Wittchen HU, Robins LN, Cottler LB, Sartorius N, Burke JD, Regier D (1991) Cross-cultural feasibility, reliability and sources of variance of the Composite International Diagnostic Interview (CIDI). The Multicentre WHO/ADAMHA Field Trials. *Br J Psychiatry* 159: 645-653, 658
9. Wittchen HU (1994) Reliability and validity studies of the WHO--Composite International Diagnostic Interview (CIDI): a critical review. *J Psychiatr Res* 28: 57-84
10. Alaghband Rad J (2003) Study of the reliability, validity, and feasibility of Farsi translation of the Composite International Diagnostic Interview (CIDI). In: Ahmadi Abhari SA, Malakooti K, Nasr Esfahani M, Razzaghi EM, Sadeghi M, Yasamy MT (eds) *Mental health effects of Iraqi invasion of Kuwait in a war – torn population of Iran: an epidemiological and financial study of the consequences of the Kuwaiti oil well fire disaster in the aftermath of Iraqi invasion of Kuwait in 1991, United Nations Compensation Commission (UNCC) Monitoring and Assessment Project*. Islamic Republic of Iran Ministry of Health, Committee for assessment and follow up for damages resulting from the Iraq–Kuwait War, Tehran - Iran,
11. Eaton WA, Chen C, Bromet EJ (2011) Epidemiology of Schizophrenia. In: Tsuang MT, Tohen MB, Jones PB (eds) *Textbook in Psychiatric Epidemiology* (3rd ed). John Wiley & Sons, New York, pp 263-287
12. Sharifi V, Assadi SM, Mohammadi MR, Amini H, Kaviani H, Semnani Y, Shabani A, Shahrivar Z, Davari-Ashtiani R, Shooshtari MH, Seddigh A, Jalali M (2009) A Persian translation of the Structured Clinical Interview for Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition: psychometric properties. *Compr Psychiatry* 50: 86-91. doi: 10.1016/j.comppsy.2008.04.004
13. Hajebi A, Motevalian A, Amin-Esmaeili M, Hefazi M, Radgoodarzi R, Rahimi-Movaghar A, Sharifi V (2012) Telephone versus face-to-face administration of the Structured Clinical Interview for Diagnostic

- 1  
2  
3  
4 and Statistical Manual of Mental Disorders, Fourth Edition, for diagnosis of psychotic disorders. *Compr*  
5 *Psychiatry* 53: 579-583.  
6  
7  
8 14. Sheehan DV, Harnett-Sheehan K, Raj BA (1996) The measurement of disability. *Int Clin*  
9 *Psychopharmacol* 11 Suppl 3: 89-95  
10  
11 15. Demyttenaere K, Bruffaerts R, Posada-Villa J et al. (2004) Prevalence, severity, and unmet need for  
12 treatment of mental disorders in the World Health Organization World Mental Health Surveys. *Jama*  
13 291: 2581-2590  
14  
15  
16 16. Druss BG, Hwang I, Petukhova M, Sampson NA, Wang PS, Kessler RC (2009) Impairment in role  
17 functioning in mental and chronic medical disorders in the United States: Results from the national  
18 comorbidity survey replication. *Mol Psychiatry* 14: 728-737  
19  
20  
21 17. Muñoz MDCL, Medina-Mora ME, Borges G, Zambrano J (2007) Social cost of mental disorders:  
22 Disability and work days lost. Results from the Mexican survey of psychiatric epidemiology. *Salud Mental*  
23 30: 4-11  
24  
25  
26 18. Amin-Esmaeili M, Motevalian SA, Rahimi-Movaghar A, Hajebi A, Hefazi M, Rad Goodarzi R, Sharifi V  
27 (2014) Sheehan disability scale: translation and psychometric assessment of the Persian version. *Iran J*  
28 *psych* 9 (In print)  
29  
30  
31 19. Hosseinpoor AR, Naghavi M, Alavian SM, Speybroeck N, Jamshidi H, Vega J (2007) Determinants of  
32 seeking needed outpatient care in Iran: Results from a national health services utilization survey. *Arch*  
33 *Iran Med* 10: 439-445  
34  
35  
36 20. Naghavi M, Jamshidi HR (2005) Utilization of Health Services. Ministry of Health, Applied Researches  
37 Secretariat, Tehran, Iran [In Persian]  
38  
39  
40 21. World Health Organization (1997) CIDI, Core Version 2.1 Trainer's Manual. World Health  
41 Organization, Geneva, Switzerland  
42  
43 22. World Health Organization (1997) CIDI, Core Version 2.1 Interview's Manual. World Health  
44 Organization, Geneva, Switzerland  
45  
46 23. Rubin DB (1987) Multiple Imputation for Nonresponse in Surveys. John Wiley and Sons, New York,  
47 NY  
48  
49 24. Bijl RV, Ravelli A (2000) Psychiatric morbidity, service use, and need for care in the general  
50 population: results of The Netherlands Mental Health Survey and Incidence Study. *Am J Public Health*  
51 90: 602-607  
52  
53 25. Henderson S, Andrews G, Hall W (2000) Australia's mental health: an overview of the general  
54 population survey. *Aust N Z J Psychiatry* 34: 197-205  
55  
56 26. Farhoudian A, Sharifi V, Amini H, Basirnia A, Mesgarpour B, Mansouri N, Amin-Esmaeili M, Salesian  
57 N, Mohammadi MR, Yousefi-Nooraie R, Rahimi-Movaghar A (2007) Prevalence of psychiatric disorders in  
58 Iran: A systematic review. *Iran J Psychiatry* 2: 137-150  
59  
60  
61  
62  
63  
64  
65

- 1  
2  
3  
4 27. World Health Organization (2011) Mental Health Atlas 2011: Country Profiles. World Health  
5 Organization, Geneva  
6  
7  
8 28. Alonso J, Angermeyer MC, Bernert S et al. (2004) Prevalence of mental disorders in Europe: results  
9 from the European Study of the Epidemiology of Mental Disorders (ESEMeD) project. *Acta Psychiatr*  
10 *Scand Suppl*: 21-27.  
11  
12 29. Jacobi F, Wittchen HU, Holting C, Hofler M, Pfister H, Muller N, Lieb R (2004) Prevalence, co-  
13 morbidity and correlates of mental disorders in the general population: results from the German Health  
14 Interview and Examination Survey (GHS). *Psychol Med* 34: 597-611  
15  
16 30. Kessler RC, McGonagle KA, Zhao S, Nelson CB, Hughes M, Eshleman S, Wittchen HU, Kendler KS  
17 (1994) Lifetime and 12-month prevalence of DSM-III-R psychiatric disorders in the United States. Results  
18 from the National Comorbidity Survey. *Arch Gen Psychiatry* 51: 8-19  
19  
20 31. Stewart P, Lips T, Lakaski C, Upshall P (2002) A Report on Mental Illnesses in Canada. Health Canada  
21 Editorial Board Mental Illnesses in Canada, Ottawa  
22  
23 32. Bromet E, Andrade LH, Hwang I et al. (2011) Cross-national epidemiology of DSM-IV major  
24 depressive episode. *BMC Med* 9: 90. doi:10.1186/1741-7015-9-90  
25  
26 33. Sadeghirad B, Haghdoost AA, Amin-Esmaeili M, Ananloo ES, Ghaeli P, Rahimi-Movaghar A, Talebian  
27 E, Pourkhandani A, Noorbala AA, Barooti E (2010) Epidemiology of major depressive disorder in iran: a  
28 systematic review and meta-analysis. *Int J Prev Med* 1: 81-91  
29  
30 34. Hawthorne G, Goldney R, Taylor AW (2008) Depression prevalence: is it really increasing? *Aust N Z J*  
31 *Psychiatry* 42: 606-616. doi:10.1080/00048670802119788  
32  
33 35. Andersen I, Thielen K, Bech P, Nygaard E, Diderichsen F (2011) Increasing prevalence of depression  
34 from 2000 to 2006. *Scand J Public Health* 39: 857-863.  
35  
36 36. Goldney RD, Eckert KA, Hawthorne G, Taylor AW (2010) Changes in the prevalence of major  
37 depression in an Australian community sample between 1998 and 2008. *Aust N Z J Psychiatry* 44: 901-  
38 910.  
39  
40 37. Wang J, Williams J, Lavorato D, Schmitz N, Dewa C, Patten SB (2010) The incidence of major  
41 depression in Canada: the National Population Health Survey. *J Affect Disord* 123: 158-163.  
42  
43 38. Noorbala AA, Bagheri Yazdi SA, Hafezi M (2012) Trends in change of mental health status in the  
44 population of Tehran between 1998 and 2007. *Arch Iran Med* 15: 201-204.  
45  
46 39. Saha S, Chant D, Welham J, McGrath J (2005) A systematic review of the prevalence of  
47 schizophrenia. *PLoS Med* 2: e141. doi: 10.1371/journal.pmed.0020141  
48  
49 40. Bijl RV, van Zessen G, Ravelli A, de Rijk C, Langendoen Y (1998) The Netherlands Mental Health  
50 Survey and Incidence Study (NEMESIS): objectives and design. *Soc Psychiatry Psychiatr Epidemiol* 33:  
51 581-586  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

41. Korten A, Henderson S (2000) The Australian national survey of mental health and well-being: Common psychological symptoms and disablement. Br J Psychiatry 177: 325-330

42. World Health Organization (2013) Global Burden of Disease Country Profile- Iran.  
<http://www.healthmetricsandevaluation.org/sites/default/files/country-profiles/GBD%20Country%20Report%20-%20Iran.pdf>. Accessed 02/02/2014 2014

**Table 1** Socio-demographic characteristics of the IranMHS respondents (N=7886)

<b>Characteristics</b>	<b>n</b>	<b>Unweighted %</b>	<b>Weighted % (SE)</b>
<b>Gender</b>			
Male	3,387	42.9	50.5 (0.7)
Female	4,499	57.1	49.5 (0.7)
<b>Age group (years)</b>			
15-19	998	12.7	18.1 (0.6)
20-29	2,549	32.3	33.8 (0.6)
30-39	2,200	27.9	21.8 (0.5)
40-49	1,188	15.0	15.3 (0.5)
50-59	704	9.0	8.7 (0.4)
60-64	247	3.1	2.5 (0.2)
<b>Marital Status</b>			
Never married	2,025	25.7	33.0 (0.7)
Married	5,527	70.1	63.5 (0.7)
Previously married	332	4.2	3.6 (0.2)
<b>Education</b>			
Illiterate	646	8.2	5.7 (0.3)
Primary school	1,917	24.4	19.6 (0.6)
Middle school	1,280	16.3	15.5 (0.5)
(High school	2,823	35.8	40.7 (0.7)
University	1,208	15.3	18.5 (0.7)
<b>Occupation</b>			
Employed	2,803	35.5	38.4 (0.7)
Students	937	11.9	17.0 (0.6)
Retired	166	2.1	2.4 (0.2)
Housewives	3,241	41.1	32.7 (0.6)
Unemployed	737	9.4	9.5 (0.4)
<b>Residence</b>			
Urban	4,380	55.5	70.9 (1.1)
Rural	3,506	44.5	29.1 (1.1)
<b>Socio-economic status</b>			
Low	2,152	28.0	22.2 (0.7)
Moderate	3,191	41.6	39.7 (0.8)
High	2,330	30.4	38.1 (0.9)

Abbreviations: IranMHS, Iranian Mental Health Survey; SE, Standard Error

**Table 2** Twelve-month prevalence of psychiatric disorders in the IranMHS (N=7886)

Psychiatric disorders	Total		Female		Male	
	n	% (95% CI)	n	% (95% CI)	n	% (95% CI)
<b>Any anxiety disorder</b>	<b>1,029</b>	<b>15.6 (14.5-16.6)</b>	<b>822</b>	<b>19.4 (17.9-20.9)</b>	<b>387</b>	<b>12.0 (10.6-13.4)</b>
Panic disorder with/ without agoraphobia	182	1.9 (1.6-2.3)	144	2.5 (2.0-3.0)	54	1.4 (0.9-1.8)
Agoraphobia without panic	118	1.5 (1.1-1.8)	91	2.0 (1.5-2.5)	27	0.9 (0.5-1.4)
Social phobia	274	3.2 (2.7-3.6)	191	4.1 (3.4-4.8)	83	2.3 (1.7-2.9)
Generalized anxiety disorder	427	5.2 (4.6-5.8)	275	5.9 (5.2-6.7)	152	4.5 (3.7-5.4)
Obsessive-compulsive disorder	408	5.1 (4.5-5.7)	297	6.8 (5.9-7.6)	111	3.4 (2.7-4.1)
Posttraumatic stress disorder	169	2.1 (1.7-2.4)	106	2.4 (1.9-3.0)	63	1.7 (1.2-2.3)
<b>Any mood disorder</b>	<b>1,153</b>	<b>14.6 (13.6-15.6)</b>	<b>766</b>	<b>17.3 (15.9-18.7)</b>	<b>387</b>	<b>11.9 (10.5-13.2)</b>
Major depressive disorder	1,014	12.7 (11.8-13.7)	678	15.4 (14.1-16.7)	336	10.2 (8.9-11.4)
Dysthymia	110	1.3 (1.0-1.5)	79	1.6 (1.2-2.1)	31	0.9 (0.5-1.3)
Bipolar I disorder	64	1.0 (0.7-1.2)	34	0.8 (0.5-1.1)	30	1.1 (0.7-1.6)
<b>Any substance use disorder*</b>	<b>185</b>	<b>2.8 (2.3-3.3)</b>		<b>0.4 (0.2-0.7)</b>		<b>5.2 (4.2-6.1)</b>
Drug abuse	30	0.5 (0.3-0.7)	1	0.01 (0.0-0.04)	29	1.0 (0.6-1.4)
Drug dependence	130	1.8 (1.5-2.2)	14	0.3 (0.1-0.5)	116	3.3 (2.6-3.9)
Alcohol abuse	15	0.3 (0.1-0.5)	0	0.0	15	0.7 (0.3-1.0)
Alcohol dependence	35	0.6 (0.4-0.9)	3	0.1 (0.0-0.02)	32	1.2 (0.7-1.6)
<b>Any primary psychotic disorder</b>	<b>31</b>	<b>0.50 (0.33-0.66)</b>	<b>16</b>	<b>0.46 (0.26-0.68)</b>	<b>15</b>	<b>0.54 (0.28-0.80)</b>
<b>Any psychiatric disorder</b>	<b>1,890</b>	<b>23.6 (22.4-24.8)</b>	<b>1,201</b>	<b>26.5 (24.9-28.2)</b>	<b>689</b>	<b>20.8 (19.1-22.4)</b>

\*Abbreviations: IranMHS, Iranian Mental Health Survey; CI, confidence Interval

\*Major categories Included: Alcohol, opioid, cannabis, stimulants, hallucinogens, and inhalants.

**Table 3** Severity proportion of 12- month psychiatric disorders in the IranMHS (N=7886)

<b>Psychiatric disorders</b>	Severity *, % (95% CI)		
	Serious	Moderate	Mild
Any anxiety disorder	30.6 (27.3-33.9)	30.1 (27.0-33.3)	39.3 (35.8-42.7)
Any mood disorder	36.0 (32.7-39.4)	33.9 (30.5-37.2)	30.1 (26.8-33.3)
Any substance use disorder **	80.6 (73.6-87.7)	7.5 (2.5-12.5)	11.9 (6.2-17.6)
Any primary psychotic disorder	100	-	-
Any psychiatric disorder	34.3 (31.6-36.9)	29.5 (27.0-32.1)	36.2 (33.6-38.8)

Abbreviations: IranMHS, Iranian Mental Health Survey; CI, confidence Interval

\* Percentages shown in each row are proportions of the cases in each group of severity and the total is equal to 100%.

\*\* Major categories Included: Alcohol, opioid, cannabis, stimulants, hallucinogens, and inhalants.



**Table 4** Socio-demographic correlates of 12-month prevalence of any psychiatric disorder in the IranMHS (N=7886)

Socio-demographic characteristics	Any psychiatric disorder	
	Unadjusted OR (95% CI)	Adjusted OR* (95% CI)
<b>Gender</b>		
Male	1	1
<b>Female</b>	1.38 (1.21-1.57)**	<b>1.47 (1.21-1.78)**</b>
<b>Age groups</b>		
15-19	1	1
<b>20-29</b>	1.20 (0.97-1.47)	<b>1.27 (0.98-1.65)</b>
30-39	1.15 (0.93-1.42)	1.19 (0.89-1.60)
40-49	1.19 (0.95-1.50)	1.21 (0.89-1.66)
50-59	1.12 (0.86-1.49)	1.08 (0.75-1.56)
60-64	0.85 (0.55-1.33)	0.75 (0.43-1.30)
<b>Marital status</b>		
Never married	1	1
<b>Married</b>	1.03 (0.89-1.19)	<b>1.07 (0.87-1.30)</b>
<b>Previously married</b>	1.91 (1.42-2.57)**	<b>1.77 (1.25-2.51)**</b>
<b>Education</b>		
Illiterate	1	1
Primary	0.97 (0.75-1.25)	0.97 (0.74-1.26)
<b>Secondary</b>	0.93 (0.71-1.22)	<b>1.02 (0.75-1.38)</b>
High school	0.85 (0.67-1.08)	0.93 (0.70-1.23)
<b>University</b>	0.70 (0.53-0.92)***	<b>0.65 (0.50-1.00)***</b>
<b>Occupation</b>		
Employed	1	1
Students	1.03 (0.83-1.27)	1.00 (0.77-1.31)
Retired	0.91 (0.58-1.41)	1.10 (0.66-1.81)
Homemakers	1.34 (1.16-1.55)**	0.90 (0.73-1.11)
<b>Unemployed</b>	1.89 (1.52-2.36)**	<b>1.81 (1.43-2.30)**</b>
<b>Residence</b>		
Urban	1	1
<b>Rural</b>	0.89 (0.77-1.02)	<b>0.71 (0.60-0.83)**</b>
<b>Socio-economic status</b>		
Low	1	1
<b>Middle</b>	0.78 (0.66-0.90)**	<b>0.74 (0.62-0.87)**</b>
<b>High</b>	0.65 (0.55-0.78)**	<b>0.64 (0.52-0.78)**</b>

Abbreviations: IranMHS, Iranian Mental Health Survey; CI, confidence Interval; OR, Odds Ratio

\* Adjusted for age, sex, marital status, occupation, education, place of residence and socio-economic status.

\*\* Statistically significant at the 0.001 level

\*\*\* Statistically significant at the 0.05 level